Rose Pharmacy 10008 Pines Blvd Pembroke Pines, FL 33024 Phone: 954.432.8290 Fax: 844.246.3364



**INFUSION ORDERS- REMICADE (INFLIXIMAB)** 

	PATIENT IN	VFORMATIO	N		
Name: DOB:					
Allergies:	Date of Re	Date of Referral:			
		<u> </u>			
	REFERRA	AL STATUS			
☐ New Referral ☐ Dose or Frequency Change ☐ Order Renewal				er Renewal	
	DIAGNOSIS A	ND ICD 10 (	ODE		
☐ Moderate to Severe Ulcerative			ICD 10 Code: K51.90		
	☐ Moderate to Severe Crohn's Disease		ICD 10 Code: K50.90		
		ICD 10 Code: M06.9			
, , ,			CD 10 Code: M45.9		
			CD 10 Code: L40.52		
1			CD 10 Code: L40.0 CD10 Code:		
Other:		ICDIO Code:			
	REQUIRED I	OOCUMENTA	TION		
☐ This signed order form by the provider		JOCOWE TVIII	☐ Clinical/Progress notes		
☐ Patient demographics AND insurance information			☐ Labs and Tests supporting primary diagnosis		
☐ Hepatitis B Test Results: HBsAg, Total HepB Core			11 31 3 3		
Antibody			☐ TB Test Results		
List Tried & Failed Therapies, incl	uding duration of treatment:				
1) 2)					
3)					
,					
	MEDICA	TION ORDERS			
Initial Dosing	☐ Remicade 5mg/kg IV at week		ry 8 weeks thereaft	er	
Maintenance Dosing	☐ Remicade 5mg/kg IV every 8 v				
Alternative Dosing	☐ Remicade	IV every	1		
TO			_weeks		
	kg	•	_weeks		
Patient Weight=  Refills: □ X 6 m		•	_weeks		
	onths	doses	_weeks		
Refills: \( \sum \text{X 6 m}	onths	•	_weeks		
Refills: ☐ X 6 m	onths	doses	_weeks		
Refills:	PREMED or to Remicade infusion or to Remicade infusion	doses	_weeks		
Refills: ☐ X 6 m  ☐ Acetaminophen 650mg PO pric ☐ Diphenhydramine 25mg PO pric ☐ Methylprednisolone 40mg Slow	PREMED or to Remicade infusion or to Remicade infusion	doses	_weeks		
Refills:	PREMED or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction	doses		s as deemed medically necessary	
Refills:	PREMED or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction ccurs, the on-call physician will or	doses  ICATIONS  rder appropriat	rescue medication	s as deemed medically necessary.	
Refills:	PREMED  or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction  ccurs, the on-call physician will or cing the rate of infusion or discont	doses  CICATIONS  rder appropriationing the median	rescue medication	s as deemed medically necessary.	
Refills:	PREMED  or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction  ccurs, the on-call physician will or cing the rate of infusion or discont	doses  ICATIONS  rder appropriat	rescue medication	s as deemed medically necessary.	
Refills:	PREMED  or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction  ccurs, the on-call physician will or cing the rate of infusion or discont	doses  CICATIONS  rder appropriationing the median	rescue medications cation.	s as deemed medically necessary.	
Refills:	PREMED  or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction  ccurs, the on-call physician will or cing the rate of infusion or discont	doses  CICATIONS  rder appropriationing the median	rescue medications ication.	s as deemed medically necessary.	
Refills:	PREMED  or to Remicade infusion or to Remicade infusion IV Push PRN infusion reaction  ccurs, the on-call physician will or cing the rate of infusion or discont	doses  CICATIONS  rder appropriationing the median	rescue medications cation.	s as deemed medically necessary.	

Contact us with questions at: info@rosenursing.net or call **954.432.8290**